

# Welche Optionen hat die Forstwirtschaft im Klimawandel?

proWALD November 2015, S. 12 ff.

## Literaturhinweise

- Beck, W. (2009): Growth patterns of forest stands - the response towards pollutants and climatic impact. *iForest* 2: 4-6 [online: 2009-01-21] URL: <http://www.sisef.it/iforest/show.php?id=472>
- Beck, W., Sanders, T., Pofahl, U. (2013): CLIMTREG: Detecting temporal changes in climate–growth reactions – A computer program using intra-annual daily and yearly moving time intervals of variable width. *Dendrochronologia* 31: 232-241
- Bolte, A., Ammer, C., Löf, M. et al. (2009): Adaptive forest management in central Europe: Climate change impacts, strategies and integrative concept. *Scandinavian Journal of Forest Research* 24: 473-482
- Brang, P., Spathelf, P., Larsen, J.B. et al. (2014): Suitability of close-to-nature silviculture for adapting temperate European forests to climate change. *Forestry* 87 (4): 492-503
- Hampe, A., Petit, R.J. (2005): Conserving biodiversity under climate change: the rear edge matters. *Ecol. Lett.* 8: 461-467
- Hanewinkel, M., Cullmann, D.A., Schelhaas, M.J. et al. (2013): Climate change may cause severe loss in the economic value of European forest land. *Nat. Clim. Chang.* 3: 203-207
- IPCC (2014): Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp.
- Jacob, D., Petersen, J., Eggert, B. et al. (2014): EURO-CORDEX: new high-resolution climate change projections for European impact research. *Regional Environmental Change*, Springer Berlin Heidelberg, 1-16
- Kätzel, R., Löffler, S. (2015): Wenn der Durst kommt – Eiche, Buche, Kiefer, Douglasie im Trockenstresstest. In: LFE (Hrsg.) *Wissenstransfer in die Praxis*. Eberswalder Forstliche Schriftenreihe, im Druck
- Lindner, M., Maroschek, M., Netherer, S. et al. (2010): Climate change impacts, adaptive capacity, and vulnerability of European forest ecosystems. *Forest Ecology and Management* 259: 698–709
- Menzel, A. (2006): Zeitliche Verschiebungen von Austrieb, Blüte, Fruchtreife und Blattverfärbung im Zuge der rezenten Klimaerwärmung. *Forum für Wissen* (Eidgenössische Forschungsanstalt WSL), 47-53
- Möller, K. (2014): Waldschutz – Management mit mehr Risiken und weniger Möglichkeiten. In: LFE (Hrsg.) *Wissenstransfer in die Praxis*. Eberswalder Forstliche Schriftenreihe, Band 55: 59-65
- Pretzsch, H., Schütze, G., Uhl, E. (2013): Resistance of European tree species to drought stress in mixed versus pure forests: evidence of stress release by inter-specific facilitation. *Plant Biol.* 15: 483-495
- Seidling, W., Ziche, D., Beck, W. (2012): Climate responses and interrelations of stem increment and crown transparency in Norway spruce, Scots pine, and common beech. *Forest Ecology and Management* 284: 196-204
- Sohn, J., Gebhardt, T., Ammer, C. et al. (2013): Mitigation of drought by thinning: Short-term and long-term effects on growth and physiological performance of Norway spruce (*Picea abies*). *Forest Ecology and Management* 308: 188-197

Spathelf, P., Bilke, G., Bolte, A. et al. (2008): Herausforderung Klimawandel – Waldmanagement im Klimastress. AFZ-DerWald 23: 1254-1255